

UNIVERSIDADE DE SÃO PAULO
INSTITUTO DE GEOCIÊNCIAS

**Geotectonic study of the southeastern Dom Feliciano Belt in southern Brazil
and its connections to the Punta del Este Terrane (Uruguay) and Pan-
African mobile belts in southwestern Africa**

RODRIGO FABIANO DA CRUZ

Doctoral Thesis presented to the Postgraduate
program of the Geoscience Institute as part of
the requirements for the degree of Doctor of
Sciences

Postgraduate Area: Igneous and Metamorphic
Petrology

Advisor: Prof. Dr. Miguel Angelo Stipp Basei

Co-advisor: Ruy Paulo Philipp

SÃO PAULO
2023

Autorizo a reprodução e divulgação total ou parcial deste trabalho, por qualquer meio convencional ou eletrônico, para fins de estudo e pesquisa, desde que citada a fonte.

Serviço de Biblioteca e Documentação do IGc/USP

Ficha catalográfica gerada automaticamente com dados fornecidos pelo(a) autor(a)
via programa desenvolvido pela Seção Técnica de Informática do ICMC/USP

Bibliotecários responsáveis pela estrutura de catalogação da publicação:
Sonia Regina Yole Guerra - CRB-8/4208 | Anderson de Santana - CRB-8/6658

Cruz, Rodrigo Fabiano

Geotectonic study of the southeastern Dom Feliciano Belt in southern Brazil and its connections to the Punta del Este Terrane (Uruguay) and Pan-African mobile belts in southwestern Africa / Rodrigo Fabiano Cruz; orientador Miguel Ângelo Stipp Basei; coorientador Ruy Paulo Philipp. -- São Paulo, 2023.

261 p.

Tese (Doutorado - Programa de Pós-Graduação em Mineralogia e Petrologia) -- Instituto de Geociências, Universidade de São Paulo, 2023.

1. U-Pb geochronology. 2. Lu-Hf isotopes. 3. Dom Feliciano Belt. 4. Western Gondwana. I. Stipp Basei, Miguel Ângelo, orient. II. Philipp, Ruy Paulo, coorient. III. Título.

UNIVERSIDADE DE SÃO PAULO
INSTITUTO DE GEOCIÊNCIAS

"Geotectonic study of the southeastern Dom Feliciano Belt in southern Brazil and its connections to the Punta del Este Terrane (Uruguay) and Pan-African mobile belts in southwestern Africa"

RODRIGO FABIANO DA CRUZ

Orientador:

Prof. Dr. Miguel Angelo Stipp Basei

Tese de Doutorado

Nº 668

COMISSÃO JULGADORA

Dr. Miguel Angelo Stipp Basei

Dr. Carlos José Archanjo

Dra. Carla Cristine Porcher

Dr. Pedro Oyhantçabal

Dra. Joseneusa Brilhante Rodrigues

SÃO PAULO
2023

*“Long is the way and hard, that out of Hell
leads up to light.”*

John Milton

AGRADECIMENTOS

Caro leitor, a presente tese de doutorado escrita como trabalho de um autor, é na realidade resultado do esforço de muitos que de uma forma ou outra, voluntariamente ou inconscientemente, contribuíram com esta jornada. Para começo de conversa, agradeço o apoio financeiro da Fundação de Amparo à Pesquisa de São Paulo (FAPESP) através do Projeto Temático número 2015/03737-0 e ao Serviço Geológico do Brasil (SGB-CPRM) pelo apoio de diversas formas, seja financeiramente, tecnicamente ou mesmo na forma da liberação para participação de disciplinas e trabalhos de campo. Sou muito grato aos meus orientadores, Miguel Basei e Ruy Philipp, pelo forte incentivo e preciosos ensinamentos.

Eternas graças aos colegas de pós-graduação da USP e colegas de trabalho da CPRM, sem citar todos pra não se esquecer de dar o devido crédito, mas com um salve especial para os que deram uma grande força nesta etapa final do doc: Lucas Lino, Lucas Tesser, Aninha, Andrea Sander, Jorge Laux, Mariana, José Luciano, Iglesias, Betiollo, Mathias e aos demais que por ventura não escrevi aqui, saibam que sempre serei muito agradecido. Expresso meu reconhecimento e gratidão as universidade públicas (UFRGS e USP), pela oportunidade de ter um ensino de qualidade e gratuito, de outra maneira seria muito, mas muito difícil chegar até aqui. Um especial salve aos professores que me deram um rumo na vida! Também faço o devido reconhecimento ao excelente trabalho da equipe técnica do CPGeo do Instituto de Geociências da USP, que geraram dados de qualidade, que em boa parte já se encontram publicados em revistas de alta qualificação pelo Qualis CAPES.

Agradeço do fundo do meu coração aos meus pais, dona Nadir e seu Romeu, ao meu irmão, minha cunhada e minhas sobrinhas, sem o apoio e carinho de vocês eu nada seria. Aos amigos que sempre que preciso estão presentes, podem vir de diferentes épocas e ambientes, dos tempos de graduação ou da firma, só quero que saibam que sou muito grato a todos (sem citar nomes, vocês sabem quem são!). Certeza que esqueci muita gente aqui, a estes não se preocupem que assim que os encontrar pessoalmente agradecerei efusivamente, mesmo que alguns provavelmente não entenderão nada ou não façam a menor ideia do que está descrito aqui. De qualquer forma, sou um afortunado por ter encontrado tantas pessoas especiais ao longo do caminho.

"In everyday life, there is more than meets the eye. To reach the depths of truth, we must drag the waters." Dimebag Darrel

RESUMO

As unidades supracrustais neoproterozoicas do Cinturão Dom Feliciano no sul do Brasil e leste uruguaio fornecem um registro da evolução tectônica do ciclo Brasileiro/Pan-Africano em termos de sedimentação, deformação, plutonismo e metamorfismo durante a assembleia do Gondwana ocidental. Foram adquiridos dados U-Pb e Lu-Hf em zircão de rochas metassedimentares e granitos do norte do Terreno Punta del Este (Domínio Jaguarão) e do Batólito Pelotas. Os complexos metassedimentares do Domínio Jaguarão possuem idades de zircão detrítico variando entre 3,2 a 0,7 Ga, com picos entre 2,2–1,7 Ga, 1,5–1,0 Ga e 1,0–0,7 Ga. As amostras do Complexo Herval, inserido no Batólito Pelotas, revelaram semelhantes idades de proveniência, com dois intervalos principais de 1,5–1,1 Ga e 1,0–0,7 Ga, com intervalos subordinados de 2,0–1,7 Ga e 2,8–2,4 Ga. A idade máxima de deposição dos complexos é restrita a populações de idade entre 700 e 650 Ma. Bordas metamórficas em zircão variam de 680 a 570 Ma. Os dados Lu-Hf dos zircões meso a paleoproterozoicas possuem valores de $\epsilon_{\text{Hf}}(t)$ entre -10 a +10 e idades modelo paleoproterozoicas a arqueanas. Os valores de $\epsilon_{\text{Hf}}(t)$ em zircões neoproterozoicos são predominantemente negativos (-15 a +5) com idades modelo meso a paleoproterozoicas. Dados U-Pb em zircão de granitos cálcio-alcalinos a peraluminosos, incluindo lentes intrusivas miloníticas nas sequências metassedimentares, apresentam idades de cristalização de 580-570 Ma, apontando para um magmatismo Ediacarano generalizado no norte do Terreno Punta del Este. Esses resultados demonstram a deposição sedimentar encerrou durante o desenvolvimento do metamorfismo e magmatismo relacionado ao arco criogeniano/ediacarano do cinturão Dom Feliciano. A escassez de rochas mesoproterozoicas na Província da Mantiqueira e áreas cratônicas próximas sugere que essas populações detríticas se originaram em rochas do sudoeste africano, provavelmente relacionados aos crátons Congo/Kalahari. Os dados U-Pb e Lu-Hf do Batólito Pelotas (arco-magmático) em zircões de rochas plutônicas com evidências de mistura e mistura de magma, bem como assimilação de rochas de embasamento, sugerem magmatismo polifásico relacionado a episódios de intenso magmatismo, potencialmente desencadeados por mudanças no ambiente tectônico. As análises U-Pb mostram idades de cristalização de xenólito de gnaiss granodiorítico em $691,5 \pm 7$ Ma, xenólito de microgranodiorito em $645,9 \pm 3,7$ Ma, monzogranito hospedeiro em $634,4 \pm 4,4$ Ma e sienogranito intrusivo em $620,1 \pm 4,7$ Ma, todos intrudidos por pegmatito sienogranítico em $60,9 \pm 1,7$ Ma. Outras análises foram realizadas em granito cálcio-alcalino (633 ± 4 Ma) e granito peraluminoso ($609,6 \pm 2,2$ Ma). Os dados Lu-Hf destacam um intervalo de idade

modelo variando de 1,7 a 1,3 Ga com valores de ϵNd entre -4 e 1, exceto para a amostra de pegmatito com idades modelo paleoproterozoicas a arqueanas (3,3-1,6 Ga) e valores de ϵNd entre -33 e 1. Os dados adquiridos enfatizam uma contribuição de fontes ligeiramente juvenis a crustais, com idades modelo variando principalmente do Estateriano-Orosiriano ao Mesoproterozóico. Os resultados apontam para um contexto geológico inicial extensional durante o Toniano (~800 Ma), com posterior deformação e metamorfismo regional ligado a estágios acrescionários e colisionais (690–570 Ma).

Palavras-chave: Geocronologia U-Pb, Isótopos Lu-Hf, Cinturão Dom Feliciano, Gondwana Ocidental

ABSTRACT

The Neoproterozoic supracrustal units of the Dom Feliciano Belt in southern Brazil and eastern Uruguay provide a record of the tectonic evolution of the Brasiliano/Pan-African cycle in terms of sedimentation, deformation, plutonism and metamorphism during the western Gondwana supercontinent assembly. From the northern Punta del Este Terrane (Jaguarão Domain) and Pelotas Batholith were acquired U-Pb and Lu-Hf zircon data from metasedimentary rocks and granites. The Jaguarão Domain metasedimentary complexes have detrital zircon ages ranging from 3.2 to 0.7 Ga, with prominent age peaks at 2.2–1.7 Ga, 1.5–1.0 Ga, and 1.0–0.7 Ga. The Herval Complex (Pelotas Batholith) revealed a similar provenance of detrital zircon ages, with two main intervals of 1.5–1.1 Ga and 1.0–0.7 Ga, and minor Paleoproterozoic (2.0–1.7 Ga) and Archean (2.8–2.4 Ga) ages. The maximum depositional age of the complexes is constrained to age populations between 700 and 650 Ma. Metamorphic zircon overgrowths range from 680 to 570 Ma. The Lu-Hf data from the meso- to Paleoproterozoic zircons have $\epsilon_{\text{Hf}}(t)$ values between -10 to +10 and model ages from Paleoproterozoic to Archean. The $\epsilon_{\text{Hf}}(t)$ values in Neoproterozoic zircons are dominantly negative (-15 to +5) with meso- to Paleoproterozoic model ages. U-Pb zircon data from calc-alkaline to peraluminous granites, including mylonitic intrusive lenses in the metasedimentary sequences, present crystallization ages of 580–570 Ma, which point to a widespread Ediacaran magmatism in the northern Punta del Este Terrane. These results demonstrate that sedimentary deposition ceased and underwent metamorphism during the development of the Cryogenian/Ediacaran arc-related magmatism of the Dom Feliciano belt. Given the scarcity of Mesoproterozoic rocks in the Mantiqueira Province and nearby cratonic areas, the obtained data suggest that these detrital populations originated in southwestern African rocks, most likely the Congo/Kalahari cratons. The Pelotas Batholith (Magmatic-arc domain) have U-Pb and Lu-Hf zircon data acquired from plutonic rocks with evidence of magma mixing and mingling, as well as assimilation of wall rocks, which suggest multi-stage magmatism related to flare-up episodes of magmatism, potentially triggered by changes in the tectonic setting. U-Pb zircon analyses yielded crystallization ages of granodioritic gneiss xenolith at 691.5 ± 7 Ma, microgranodiorite xenolith at 645.9 ± 3.7 Ma, host monzogranite at 634.4 ± 4.4 Ma, and intrusive syenogranite at 620.1 ± 4.7 Ma, all intruded by syenogranitic pegmatite at 606.9 ± 1.7 Ma. Other analyses were performed on calc-alkaline granite (633 ± 4 Ma) and peraluminous granite (609.6 ± 2.2 Ma). The Lu-Hf data highlights a model age interval ranging from 1.7 to 1.3 Ga with ϵ_{Nd} values between -4 and 1, except for the pegmatite

sample with Paleoproterozoic and Archean model ages (3.3-1.6 Ga) and ϵNd values between -33 and -1. Furthermore, the acquired data emphasizes a contribution from slightly juvenile to crustal sources, with model ages ranging primarily from Statherian-Orosirian to Mesoproterozoic. The geological context and isotopic results point to a prior extensional tectonic setting during the Tonian (~800 Ma), with regional deformation and metamorphism linked with accretionary and collisional stages (690–570 Ma).

Keywords: U-Pb geochronology, Lu-Hf isotopes, Dom Feliciano Belt, Western Gondwana.

SUMMARY

1.	INTRODUCTION.....	13
1.1	Motivation for research and overarching goals.....	13
1.2	Dissertation outline.....	16
2.	MATERIAL AND METHODS.....	17
2.1	Field work and general aspects of the geological units.....	17
2.2	Microscopic Petrography and whole-rock chemical analysis.....	18
2.3	Geochronological and isotopic analysis of rocks.....	18
2.4	Isotope geology.....	18
2.4.1	The U-Pb method and zircon	19
2.4.2	The Lu-Hf Method in zircon.....	20
2.4.3	Geochronological data acquisition equipment.....	20
2.4.4	Zircon imaging techniques.....	21
2.4.5	Detrital zircon provenance studies.....	21
3.	UNVEILING THE STRATIGRAPHY OF THE NORTHERN PUNTA DEL ESTE TERRANE (JAGUARÃO DOMAIN) AND ITS ROLE IN THE NEOPROTEROZOIC TECTONIC EVOLUTION OF THE DOM FELICIANO BELT (SOUTHERN BRAZIL AND URUGUAY).....	23
3.1	Introduction.....	24
3.2	Geotectonic setting.....	25
3.2.1	Geology of the northern Punta del Este Terrane (Jaguarão Domain).....	27
3.3	Methodology.....	35
3.3.1	Geophysical data.....	35
3.3.2	Whole-rock chemistry.....	36
3.3.3	U-Pb zircon data.....	36
3.4	Results.....	38
3.4.1	Structural Data.....	38
3.4.2	Geophysical Data.....	39
3.4.3	Whole-rock chemical analyzes.....	40
3.4.4	U-Pb zircon geochronology.....	42
3.5	Discussions.....	43
3.5.1	Geophysical data.....	43
3.5.2	Granitic Magmatism.....	44
3.5.3	U-Pb provenance data.....	45

3.5.4	Tectonic significance.....	48
3.6	Conclusions.....	51
3.7	References.....	51
3.8	APPENDIX A – supplementary material of chapter 3.....	63
4.	INSIGHTS INTO THE EVOLUTION OF THE SOUTHEASTERN DOM FELICIANO BELT AND ITS CONNECTION TO THE PAN-AFRICAN OROGENY BASED ON NEW U-PB AND LU-HF ZIRCON DATA.....	74
4.1	Introduction.....	75
4.2	Geotectonic setting.....	76
4.2.1	Geology of the Jaguarão Domain/Northern Punta del Este Terrane.....	79
4.3	Methodology.....	82
4.4	Results.....	83
4.4.1	U-Pb geochronology.....	83
4.4.2	Lu-Hf zircon analyses.....	91
4.5	Discussion.....	92
4.5.1	U-Pb results, maximum depositional ages and metamorphism.....	92
4.5.2	Potential sources areas.....	95
4.5.3	Correlation of U-Pb ages in detrital zircon with Pan-African and Brasiliano metasedimentary complexes and foreland basins.....	97
4.5.4	Lu-Hf results and comparisons with the units from Dom Feliciano Belt and Pan- African Orogeny.....	101
4.5.5	Tectonic significance.....	103
4.6	Conclusions.....	105
4.7	References.....	106
4.8	APPENDIX A – supplementary material of chapter 4.....	120
	References.....	124
5.	POTENTIAL FLARE-UPS AND LULLS IN THE MULTI-STAGE MAGMATISM OF THE DOM FELICIANO BELT, SOUTHERN BRAZIL: EVIDENCE FROM GEOCHEMISTRY AND ISOTOPIC DATA.....	165
5.1	Introduction.....	166
5.2	Geotectonic setting.....	169
5.2.1	Pelotas Batholith.....	171
5.2.2	Northern Punta del Este Terrane-Jaguarão Domain.....	175
5.3	Methodology.....	175

5.4	Results.....	176
5.4.1	Chemical features of the magmatic suites.....	176
5.4.2	U-Pb geochronology and Hf isotopes.....	181
5.5	Discussion.....	188
5.5.1	Possible flare-ups and lulls/steady-state events.....	188
5.5.2	Tectonic setting models.....	194
5.5.3	The framework and generation of the Pelotas Batholith.....	196
5.6	Conclusions.....	199
5.7	References.....	200
5.8	APPENDIX A – supplementary material of chapter 5.....	210
	Whole-rock chemical analysis.....	215
	References.....	215
6.	CONCLUSIONS AND REMARKS	253
7.	REFERENCES.....	255

1. INTRODUCTION

1.1 Motivation for research and overarching goals

Studies on supercontinent tectonic settings and the evolution of Precambrian terranes have made significant progress in recent decades due to the contribution of geophysical and geochronological data through advances in the determinations of U-Pb crystallization ages and Lu-Hf isotopic composition in zircons, allowing a more accurate definition of geological domains. (Andersen, 2005, Condie, 2011; Cawood et al., 2012). The Neoproterozoic supracrustal units of the Dom Feliciano Belt (DFB) in southeastern South American Platform provide a record of the tectonic evolution of the Brasiliano/Pan-African cycle in terms of sedimentation, deformation, and metamorphism during the Western Gondwana supercontinent assembly. Indeed, the tectonic study of the DFB, which spans southwestern Brazil and eastern Uruguay over a range of ca. 1200 km (Basei et al., 2005; Philipp et al., 2016), is essential for understanding the stages of Precambrian supercontinents evolution.

The Western Gondwana (Figura 1.1) is result of the continental assembly during the Brasiliano and Pan-African cycles, whose records are found in Precambrian terranes from South America and southeastern Africa (Dalziel, 1997; Basei et al., 2005; Silva et al., 2005; Philipp et al., 2016; Hueck et al., 2018). Most paleogeographic reconstructions of Western Gondwana converge on the hypothesis that its formation process began with the breakup of Rodinia (ca. 840–750 Ma) and subsequent convergence of continental blocks (ca. 720-550 Ma), which included the Rio de La Plata, Luis Alves, Kalahari, and São Francisco-Congo cratons (Basei et al., 2008; Oyhantçabal et al., 2009; Saalman eta al., 2010; Frimmel, 2018; Hueck et al., 2018). While this general evolution is well accepted, many details are still controversial, such as subduction direction and origin of some cratonic blocks (Frimmel et al., 2011; Oyhantçabal et al., 2011; Chemale, 2012; Basei et al., 2018; Oriolo et al., 2019; De Toni et al., 2020; Hueck et al., 2022; Silva Lara et al., 2022). However, recent studies have challenged this tectonic setting as a whole, revisiting the prior concept of an intracontinental orogen (Torquato and Cordani, 1981; Porada, 1989). These studies support the model that Brasiliano/Pan-African basins developed in an intracontinental setting and subsequently underwent inversion, leading to the formation of collisional belts between ca. 650-600 Ma, followed by transcurrent tectonics between ca. 600-560 Ma. (Meira et al., 2015; Konopásek et al., 2020; Percival et al., 2021).

The idea of the present study arose during the geological mapping activity of the Sudeste do Rio Grande do Sul project (Cruz, 2019) in an area that has historically been

described as part of the Pelotas Batholith, an extensive granite massif that extends across the eastern part of the DFB. However, some previous works have pointed to the presence of a “new” tectonic block in the southern segment of the batholith. CPRM geophysical maps (CPRM, 2010) show an extensive magnetic lineament with ENE-WSW orientation that separates two distinct geophysical domains in this sector. Ramos and Koester (2014) based on geophysical maps and studies of metaultramafic rocks, highlight the area as likely block boundary.

In a mapping report for the Curral de Pedra and Passo São Diogo maps (Iglesias et al., 2018), it was concluded that part of the area covered by the projects is a different geological domain, which they termed Jaguarão Terrane. The area contains a sequence of metasedimentary rocks that form a lithological association very different from that found in the Pelotas Batholith and very similar to those found in the Punta del Este Terrane (Preciozzi et al., 1999; Basei et al., 2005), another segment of the DFB.

The primary purpose is to characterize the boundary and relation between the Pelotas Batholith and northern Punta del Este Terrane, in this research referred to as the Jaguarão Domain (Fig. 1.1). With emphasis on newly acquired isotopic data, supported by field work, thin section description, along with interpretation of whole-rock chemistry and geophysical maps. This study aimed at exploring through U-Pb and Lu-Hf zircon analyses the provenance sources of the supracrustal metamorphic complexes of the Jaguarão Domain. Additionally, to characterize the regional metamorphism and related magmatism on the domain were also acquired isotopic data from mylonitic leucogranites and porphyritic plutons that crosscut these complexes.

Futhermore, new zircon U-Pb and Lu-Hf SHRIMP/LA-ICPMS data were obtained in granitoids of the Pinheiro Machado Suite, the largest unit of the Pelotas Batholith, to provide new insights or refine existing knowledge about the magmatism of DFB. The suite exhibits evidence of magma recharges, assimilation of xenoliths, mingling, and mixing with mafic magmas, as observed in previous studies (Philipp et al., 2002; Philipp and Machado, 2005; Loureiro et al., 2021). Along with the isotopic data, interpretation of the whole-rock chemistry was used to support the findings.




Figure 1.1: A) Reconstruction of the South American and African continents. B) Former Western Gondwana terranes framework with outlined study area (modified from Basei *et al*, 2008, Goscombe *et al*, 2018, Frimmel, 2018, Hueck *et al*, 2018). Shear zones (SZ): IPSZ - Itajaí-Perimbó; MGSZ - Major Gercino; ISZ - Ibaré Shear Zone. DCSZ - Dorsal do Canguçu; SBSZ - Sierra Ballena; TPSZ - Three Palms; PSZ – Purros; ST - Sesfontein Thrust; KGT - Khorixas-Gaseneirob Thrust; OT – Otjorongo Thrust; HFT - Hakos Frontal Thrust; SKT – Schakalsberge Thrust; PWT - Piketberg-Wellington Thrust; BRT – Blackridge Thrust.

1.2 Dissertation outline

This doctoral dissertation has been organized into six main chapters. Chapter 1 introduces the motivation, objectives and implications, as well as the geotectonic context of the studied area and presents the overall structure of the dissertation. Chapter 2 includes a brief review of the applied methodology of the proposed research. Chapters 3, 4, and 5 present the results obtained during the research, which were organized and included in three manuscripts of scientific papers. The first manuscript is under review by the International Geology Review journal from Taylor & Francis editorial company, and the other two are currently available in Precambrian Research and Journal of South American Earth Sciences from Elsevier academic publishing company.

Chapter 3 presents the paper “Unveiling the stratigraphy of the northern Ponta del Este Terrane (Jaguarão Domain) and its role in the Neoproterozoic tectonic evolution of the Dom Feliciano Belt (southern Brazil and Uruguay)”. The manuscript describes the basic stratigraphy of the so-called Jaguarão Domain through outcrops and thin section descriptions. The understanding and drawing of regional structures are accomplished by structural analysis based on fieldwork and geophysical maps (magnetometric and gravimetric). Additionally, LA-ICPMS U-Pb zircon and whole-rock chemical analyses were carried out to characterize the widespread peraluminous magmatism in the domain.

The paper “Insights into the evolution of the southeastern Dom Feliciano Belt and its connection to the Pan-African Orogeny based on new U-Pb and Lu-Hf zircon data” is in Chapter 4. The manuscript presents a set of results from LA-ICPMS U-Pb and Lu-Hf in zircon, including provenance analyses of metasedimentary sequences and crystallization ages of the granites that occur as intrusive lenses or plutons. To understand the geodynamics of the Jaguarão Domain are carried out Hf isotopic studies towards the most representative U-Pb zircon ages. All results have been included in the dataset of the Pan-African/Brasiliano belts from southeastern Brazil, eastern Uruguay and southwestern Africa to make the correlations and interpretations on the tectonic events that involved the domain in the context of Western Gondwana assembly.

Chapter 5 carry the paper intituled “Potential flare-ups and lulls in the multi-stage magmatism of the Dom Feliciano Belt, southern Brazil: Evidence from geochemistry and isotopic data”. The last manuscript brings U-Pb and Lu-Hf zircon analyses by SHRIMP and LA-ICPMS methods obtained primarily on samples from a single outcrop with features of magma mixing, assimilation of wall-rock, and late intrusive magmatism. The new isotopic results were compared with previously published data from the Pelotas Batholith, which,

together with whole-rock chemistry from SGB-CPRM (Brazilian Geological Survey), allows for the inference of potential events of intense magmatism associated with changes in the Dom Feliciano Belt then-ongoing tectonism.

Finally, Chapter 6 summarizes the results and discussions from the former chapters in a brief integrative conclusion of the dissertation. Furthermore, the remaining questions have been identified as potential future research topics. Emphasizing gaps and alternative techniques to generate strong outcomes for assessing the tectonic evolution of the Dom Feliciano Belt through the Western Gondwana assembly processes.

2. MATERIAL AND METHODS

2.1 Field work and general aspects of the geological units

The results presented in the papers (Chapters 3 to 5) are based on the problems identified in the mapping work of the Sudeste do Rio Grande do Sul project (Cruz, 2019), in which, supported by the interpretation of aeromagnetic and gravimetric data, the high-deformation zones are traced, as well as the definition of the central units of the Jaguarão Domain and the southern Pelotas Batholith. Subsequently, the most representative outcrops were selected to collect material for petrographic, chemical and geochronological analysis. The fieldwork, accompanied by advisors, comprises the description of 55 outcrops focusing on the following key units of the study area:

- The Arroio Telho and Arroio Grande metavolcanosedimentary complexes were describe in terms of deformation phases, metamorphism and correlations with the La Micaela Schists and La Tuna Serpentinities from Paso del Dragon (Uruguay).
- The Herval Complex, mainly composed of quartzites and subordinately of schists and metaplutonic rocks, is understood as a mega-xenolith within the Pelotas Batholith and was selected to investigate its correlation with the above complexes.
- The Três Figueiras Suite, which includes intrusive peraluminous granites, is mainly located at the boundary of the Jaguarão Domain and the Pelotas Batholith. Additionally, the Capão do Leão, Chasqueiro and Bretanha granites were also described with the objective of understanding the latter's magmatism and deformation in the Jaguarão Domain.
- The Pinheiro Machado Suite, the largest suite of the Pelotas Batholith, focused on the Prainha dos Melões outcrop, which shows a complex relationship between xenolith assimilation and tardi-intrusive magmatism, which could be understood as a fractal of the

magmatic processes in the batholith as a whole.

2.2 Microscopic Petrography and whole-rock chemical analysis

The researcher used the facilities of the SGB-CPRM (Laboratory of Porto Alegre) to prepare the 150 thin sections, and carried out the prior treatment of the 160 samples for whole-rock chemical analysis, which involved fragmenting and pulverizing the samples to achieve a particle size less than 200 mesh. The petrographic analyses of the thin sections allowed the identification of the protoliths based on the recognition of primary structures, textures and the characterization of microstructures and metamorphic conditions.

The multi-elemental analyses were carried out at SGS GEOSOL - Laboratory LTDA, (Vespasiano, Brazil). The method applied included lithium borate fusion and determination by X-ray fluorescence for major oxides and ICP-MS for minor elements and REEs (Rare Earth Elements). The acquisition of whole-rock data had the initial goal of supporting the SGB-CPRM projects from [Cruz, 2019](#), and [Takehara and Laux, 2019](#). The geochemical diagrams were created using the GCDkit program ([Janoušek et al., 2006](#)), using the whole-rock chemistry data above quoted in addition to data gathered from academic papers from [Klein et al., 2018](#), [Vieira et al., 2016](#) and [Silva et al., 2016](#). The whole-rock chemical analysis tables are presented in the Appendix of Chapters 3 and 5.

2.3 Geochronological and isotopic analysis of rocks

Once the sampling sites were selected, the work of sample preparation and isotopic analysis was carried out. Details of the procedures used for both preparation and analysis are given in the appendices to the papers in Chapters 3 to 5. Zircon age determinations were performed on 6 samples from the Arroio Telho, Arroio Grande and Herval complexes. For the granitoids, 12 U-Pb geochronologic analyses were performed. Lu-Hf isotopic studies were carried out on 4 samples of metasediments and 7 samples of granitoids. A brief bibliographical background of the geochronological studies used in this thesis is given hereafter.

2.4 Isotope geology

Estimates of the Earth's age and its rocks have gone through a long and contentious historical process, with attempts to obtain ages from rocks, such as that of James Hutton and collaborators in the 18th century, who arrived at an estimate of 700 Ma as the time elapsed since the Cambrian by calculating the sedimentation rate of different types of strata. With the discovery of radioactivity by [Becquerel \(1896\)](#), significant progress in understanding and applying the phenomenon as a technique for dating rocks began, with [Rutherford \(1905\)](#) producing the first acknowledged result by examining the decay of the element Radium to

Helium. Since then, a number of techniques in the field of isotope geology have been developed as a chronological tool, including the Rb-Sr and Sm-Nd methods, among others, but with a strong emphasis on the U-Pb method.

2.4.1 The U-Pb method and zircon

The U-Pb method is the most robust geochronological tool and is based on the principle of radioactive decay of the U, Th, and Pb isotopes. Isotopes are elements that have the same atomic number (protons) but a different mass number. Radiogenic isotopes have a decay time when they lose half their mass and transition from a parent element to a daughter element, and this elapsed time is known as their half-life ($T_{1/2}$). Zircon ($ZrSiO_4$) crystallizes with a high U concentration, has low Pb diffusion, and is incompatible with common lead. It is a widely distributed accessory mineral in the earth's crust, including igneous, sedimentary, and metamorphic rocks, as well as meteorites and lunar rocks. Zircon is chemically resilient and refractory, withstanding weathering, transport, diagenesis, and, to a lesser extent, metamorphism. Because of these properties, zircon is the primary material to employ the U-Pb technique (Hanchar & Miller, 1993; Dickinson, 1995).

The decay of U into two series of Pb isotopes produces three separate geochronometers, $^{238}U/^{206}Pb$, $^{235}U/^{207}Pb$, and $^{207}Pb/^{206}Pb$, which should all yield the same age value for the material analyzed. Wetherill (1956) proposed the concord diagram, which puts these ages into a single diagram and makes it easy to comprehend potentially discordant data. The curve of the diagram is defined by systems with concordant U-Pb ages. The isotopic compositions of the items to be studied are thus compared with the concordance curve, allowing their age to be estimated. Geochronometers generate concordant ages when the mineral to be dated has stayed closed to U loss and the corrections for Pb initially integrated into the mineral are clearly specified, and the results can align on a straight line or a route of Pb loss, known as a discordia line (Faure 1986).

The Concordia Age (Ludwig, 1998) exists when the ratios $^{207}Pb/^{235}U$, $^{206}Pb/^{238}U$, and $^{207}Pb/^{206}Pb$ are alike within uncertainty estimates, or when the corresponding U/Pb and Pb/Pb ratios have error ellipses that overlap the Concordia curve within the 95% certainty estimate (e.g. Fig. 2.1). The Tera-Wassenburg diagram (Wendt, 1984) is another type of diagram for displaying isotopic data (e.g. Fig. 2.1). The graphic was designed to make it easier to obtain data from younger rocks. Where the abscissa is the $^{238}U/^{206}Pb$ ratio, and the ordinate is the $^{207}Pb/^{206}Pb$ ratio. The individual findings that form a cohesive grouping are organized along a simple mixing line (with the common Pb) in this concordance, and the age is derived by crossing the mixing line (discordia) with the abscissa. This form of graphic depicts the lower

intercept age and has the main advantage of allowing the identification of common ^{204}Pb correction when measured in little amounts (Parrish & Noble, 2003).




Figure 2.1: A) Wetherill Concordia diagram. B) Tera-Wasserburg Concordia diagram. The diagrams used as examples are found within Chapter 04.

2.4.2 The Lu-Hf Method in zircon

In addition to the U-Pb ages obtained in zircon, hafnium (Hf) isotope measurements can be obtained, which indicate the relative composition of the source of the mineral, whether there was a greater contribution of material of mantle origin (juvenile) or material from recycled continental crust (Harrison et al. 2005). Hf isotopic values are used in a calculation called Hafnium Epsilon (ϵHf), which takes into account the known average value of the element in the CHUR (Condritic Uniform Reservoir), assuming that positive values correspond to mantle-derived magmas and negative values correspond to crustal magma-derived rocks.

2.4.3 Geochronological data acquisition equipment

The determination of the ratio of radioactive isotopes and radiogenic isotopes involves the separation of isotopes according to their mass. Mass spectrometry originated in 1913 with the development of the first magnetic field-based spectrometer (Dickin, 1995). Thermal ionization, plasma induction, and ion beam source are three widely used methods in isotope geology and geochronology. LA-ICP-MS (laser ablation inductively coupled plasma mass spectrometry) and SHRIMP (sensitive high-resolution ion microprobe) are the most regularly used spectrometers today. Both provide ages for a specific zone of the grain with high analytical accuracy and are widely used for U-Pb and Lu-Hf ages in zircon and monazite minerals. In general, these mass spectrometers consist of an energy emitter that extracts

material from the sample and then passes through an electrostatic analyzer and magnetometer system where the isotopes that reach the multicolectors are separated and ratio readings are taken.

2.4.4 Zircon imaging techniques

The interpretation of the analytical results obtained by SHRIMP and/or LA-ICP-MS techniques involves the observation of the structures and internal characteristics of the minerals analyzed, aided by images obtained by cathodoluminescence (CL) or scanning electron microscope (SEM) (Rubatto & Gebauer, 1998; Silva, 2005). Image analysis enables the recognition of different domains based on the textural pattern presented by the zircon, avoiding heterogeneous and fractured zones within the grain, thereby improving control over the acquisition and interpretation of geochronological data. Cathodoluminescence images indicate the relative uranium concentration of zircon by luminescence intensity (Rubatto & Gebauer, 1998). Magmatic textures are characterized by oscillatory zoning caused by alternating U-rich and Pb-poor halos of high and low luminescence, respectively. Metamorphic zircons generally lack oscillatory zoning and exhibit a homogeneous internal texture.

2.4.5 Detrital zircon provenance studies

U-Pb dating of detrital zircons from siliciclastic metasedimentary rocks is a widely used tool for the geological reconstruction of metamorphic terrains that have undergone at least one or two tectonic events, helping to understand the geological evolution of mobile belts and paleocontinents (Froude et al., 1983; Condie et al., 2009). The LA-ICP-MS is the most suitable equipment for detrital zircon provenance studies because it allows the spot analysis of a large number of zircons in a relatively short time (an average of 20 grains per hour session). Dating a significant number of zircons ensures that a proper provenance analysis is carried out. According to Vermeesch (2004), a value greater than 100 grains is appropriate so that no fraction containing at least 5% of the total sample population goes undetected, thereby achieving a 95% confidence level. Provenance ages are critical for understanding crustal evolutionary processes (Harrison et al., 2005).

Whole-rock Sm-Nd analyses are useful in these investigations, but often yield mixed ages, leading to misinterpretation. In these circumstances, combined U-Pb and Lu-Hf isotopic analyses in detrital zircons allow the distinction of grains of the same crystallization age but derived from crustal areas of different mantle-derived ages. As a result, it is possible to construct graphs that plot the ϵ_{Hf} parameter on the ordinate against the crystallization age on the abscissa as an indication of domains of different crustal residence ages in a particular

section of the crust.

Patterns of age signatures of detrital zircons generally reflect the tectonic setting of the basin in which they were deposited. Convergent plate margins (accretionary tectonics) are characterized by large proportions of zircons with ages close to the basin depositional interval, in this case the forearc, backarc and trench zones (Condie et al., 2009). Whereas basins formed from continental and extensive collisional environments contain higher proportions of zircons with ages older than the time of deposition, reflecting the evolutionary history of the bedrock on which they were formed (Cawood et al., 2012, 2013).

However, magmatism volume and preservation potential vary significantly with tectonic setting (Hawkesworth et al., 2009), affecting the amount of zircon present in the sedimentary record. In convergent plate settings, large volumes of magma are generated, but with low preservation potential in the geologic record (Scholl and Von Huene, 2009). Continental collision-dominated settings produce small magmatic volumes, which are melting products of pre-existing crust (Hawkesworth et al., 2010), but their preservation in the geologic record is favored during the supercontinent formation cycle, resulting in a larger amount of preserved zircons. Extensive settings generate variable amounts of magma (White, 1992), but with a greater occurrence of mafic magmatism (Storey, 1995), resulting in a more restricted generation of zircons. Furthermore, these rocks are more susceptible to erosion, resulting in reduced preservation potential. Thus, the magmatic age record is dominated by periods of final supercontinent formation, not because of massive crustal generation, but because it provides a suitable setting for the preservation of rocks generated by continental collisional magmatism.

3. CONCLUSIONS AND REMARKS

The Jaguarão Domain (JD) is the northern segment of the Punta del Este Terrane (PET), which occurs on the eastern edge of the DFB in contact with the Aiguá and Pelotas batholiths (650-550 Ma). Gravimetric geophysical maps distinguish the JD as a high density domain compared to the granitic batholith areas. It is also observed in the geophysical (aeromagnetic) maps that the main structures (Pedro Osório, Arroio Grande and Cerro Amaro shear zones) partially mark the boundaries of the JD.

Provenance data for the JD metasedimentary units showed approximately 80% zircon crystals with Meso- to Paleoproterozoic ages, distinguishing these units from similar rocks found in the DFB where Neo- and Paleoproterozoic ages predominate (e.g., Porongos and Lavallega complexes). Geologic similarities and zircon provenance patterns strongly support the correlation of the Arroio Telho/Arroio Grande and Herval complexes of Brazil with the Paso del Dragon/Cerro Olivo complexes of Uruguay and the metasedimentary sequences of southeastern Africa.

The Lu/Hf isotopic data from JD present model ages between 3.6 – 1.0 Ga and ϵ_{Hf} values predominantly around -10 to +10, which implies mixing sources originated mainly from the assembly processes of the Columbia (2200-1500 Ma) and Rodinia (1400-950 Ma) supercontinents. Tectonic and metamorphic similarities, plus the zircon provenance pattern and isotopic data, support the correlation of the JD/PET with the western domains of Gariep and Kaoko belts units (Africa).

The data set of crystallization and provenance ages places the JD in a more reliable setting as a rift-related basin on the western border of the Namaqua-Natal Belt that underwent a tectonic inversion after the collision (~660 Ma) between the African and South American domains. The metamorphic conditions reached a peak around 650-630 Ma. Then, around 570 Ma, a second thermal-metamorphic event promoted partial melting of the metasedimentary pile, resulting in widespread peraluminous magmatism in the JD and southernmost Pelotas Batholith (PB).

The emplacement of the DFB granitic batholiths marked an important change in the provenance age scenario. The input of late-Mesoproterozoic and Tonian material disappears in syn- to post-orogenic sequences of the western branch of the DFB (e.g., upper sequence of Porongos Complex, Arroyo del Soldado, and Camaquã basins). In contrast, the eastern sequences (e.g., Rocha Formation) show that the provenance did not end after the accretionary-collisional setting. The breakup of the Pangea supercontinent resulted in the current JD arrangement, located in southeastern South America as a remnant of the orogenic

systems found in southwestern Africa.

The Prainha dos Melões outcrop is a fractal evolution of the Pinheiro Machado Suite and the PB magmatism as a whole. The differentiation processes assert the influence of partially assimilated older rocks and mafic magmas through new magmatic pulses. There is also recharge of the magma chamber in the form of dykes and sills. Whole-rock chemistry shows that southeastern DFB magmatic suites are primarily classified as I-Type peraluminous granites, with some suites being S-Type peraluminous. The chemistry diagrams depict magmatism with a diversity of sources and orogenic signatures, primarily related to a collisional to post-collisional setting with a major contribution from crustal sources.

The U-Pb zircon ages support a scenario in which the magmatism evolved in pulses around 690 Ma, 645 Ma, 635 Ma, 620 Ma, and 610 Ma. The Hf isotopes from the same samples show $T_{DM}(t)$ ages with the main interval of 1.7-1.3 Ga and ϵ_{Hf} values ranging from 1 to -4. These results are similar to those found in the metasediment data, indicating that African crust had an important contribution to the isotopic signature of arc-magmatism. The gathered U-Pb dataset shows well-defined intervals of magmatic pulses, lasting on average 10 Ma, in 660-650 Ma, 640-630 Ma, 620-610 Ma, 600-590 Ma, and 580-560 Ma. The main peak of the U-Pb ages falls within the 620-610 Ma interval, which overlaps the average age of the intrusive peraluminous granites in the western boundary of the PB and is associated with the partial melting of the Várzea do Capivarita Complex.

Based on the obtained results and conclusions, I made the following recommendations:

- The development of thermodynamic modeling, which can be done by carrying out whole-rock chemistry in combination with chemical analysis of the minerals in the metamorphic paragenesis of the metasedimentary complexes, in order to generate pseudo-sections to quantify the P (pressure) and T (temperature) parameters of metamorphism.
- Perform a refined processing of the geophysical data to generate products such as magnetic deconvolution and gravimetric inversion profiles, to better delineate geophysical anomalies, as well as the depth and angle of regional structures, related to tectonic block boundaries.
- Increase the coverage of Lu-Hf data on the magmatic suites of the batholith. There is a reasonable amount of Sm-Nd data in the literature, but as they are isotopic results on whole-rock it limits more refined interpretations on the sources and evolution of magmatism.

- Conduct a refined isotopic study on the rocks of the Matarazzo Complex, another mega-xenolith of metasedimentary rocks within plutonic rocks, in order to improve the correlations of these rocks with those present in this study.

4. REFERENCES

- Andersen, T., 2005. Detrital zircons as tracers of sedimentary provenance: limiting conditions from statistics and numerical simulation. *Chem. Geol.* 216, 249–270.
- Basei, M. A. S., Frimmel, H. E., Nutman, A. P., Preciozzi, F., & Jacob, J., 2005. A connection between the Neoproterozoic Dom Feliciano (Brazil/Uruguay) and Gariep (Namibia/South Africa) orogenic belts - Evidence from a reconnaissance provenance study. *Precambrian Research*, 139(3–4), 195–221.
- Basei, M. A. S., Frimmel, H. E., Nutman, A. P., & Preciozzi, F., 2008. West Gondwana amalgamation based on detrital zircon ages from Neoproterozoic Ribeira and Dom Feliciano belts of South America and comparison with coeval sequences from SW Africa. *Geological Society, London, Special Publications*, 294(1), 239–256.
- Basei, M.A.S., Frimmel, H.E. Campos Neto, M.C., Araujo, C.E.G., Castro, N. A., Passarelli, C. R., 2018. The Tectonic History of the Southern Adamastor Ocean Based on a Correlation of the Kaoko and Dom Feliciano Belts. In: Siegesmund, S., Basei, M. A. S., Oyhantçabal, P., Oriolo, S. (eds), *Geology of Southwest Gondwana*, pages 63-85. Springer International Publishing.
- Becquerel, H., 1896. Sur les radiations invisibles emises par phosphorescence. *Compt. Rend. Acad. Sci. Paris.*, 122: 420-421.
- Cawood, P.A., Hawkesworth, C.J., Dhuime, B., 2012. Detrital zircon record and tectonic setting. *Geology*, 40 (10), 875–878.
- Cawood, P.A., Hawkesworth, C.J., Dhuime, B., 2013. The continental record and the generation of continental crust. *Geological Society of America Bulletin* 125 (1–2), 14–32.

- Chemale, F., Mallmann, G., Bitencourt, M.F., Kawashita, and K., 2012, Time constraints on magmatism along the Major Gercino Shear Zone, southern Brazil: implications for West Gondwana reconstruction. *Gondwana Res.* v.22(1), p.184-199.
- Condie, K.C., Belousova, E., Griffin, W.L., Sircombe, K.N., 2009. Granitoid events in space and time: Constraints from igneous and detrital zircon age spectra. *Gondwana Research* 15, 228-242.
- Condie, K.C., 2011. *Earth as an Evolving Planetary System*. 2nd ed. Amsterdam, Elsevier, 574p.
- CPRM - SERVIÇO GEOLÓGICO DO BRASIL. Projeto Aerogeofísico Escudo do Rio Grande do Sul: relatório final do levantamento e processamento dos dados magnetométricos e gamaespectrométricos. Rio de Janeiro: Lasa Prospecções, 2010.
- Cruz, R.F., 2019. Projeto Sudeste do Rio Grande do Sul: escalas 1:250.000 e 1:100.000; estado do Rio Grande do Sul / Relatório do Programa geologia, mineração e transformação mineral. CPRM, Porto Alegre.
- Dalziel, I.W.D., 1997. Neoproterozoic/Paleozoic geography and tectonics: Review, hypothesis, environmental speculation, *Geol. Soc. Am. Bull.* 108, 16-42.
- De Toni, G.B., Bitencourt, M.F., Nardi, L.V.S., Florisbal, L.M., Almeida, B.S., Geraldés, M., 2020. Dom Feliciano Belt orogenic cycle tracked by its pre-collisional magmatism: The Tonian (ca. 800 Ma) Porto Belo Complex and its correlations in southern Brazil and Uruguay. *Precambrian Res.* 342, 105702.
- Dickinson, A., 1995. *Radiogenic Isotope Geology*: Cambridge University Press, 490 p.
- Dodson, M.H., Compston, W., Williams, I.S., Wilson, J.F., 1988. A search for ancient detrital zircons in Zimbabwean sediments. *Geological Society (London) Journal* 145, 977-983.
- Faure, G., 1986. *Principles of Isotope Geology*. John Wiley e Sons Inc. Second Edition. 589p.

- Frimmel, H.E., 2018. The Gariep Belt. In: Siegesmund, S., Basei, M.A.S., Oyhantçabal, P., Oriolo, S. (eds), *Geology of Southwest Gondwana*, pages 353-386. Springer International Publishing.
- Froude, D.O., Ireland, T.R., Kinney, P.D., Williams, I.S., Compston, W., 1983. Ion microprobe identification of 4,100– 4,200 Myr-old terrestrial zircons. *Nature* 304, 616–618.
- Goscombe, B. Foster, D.A., Gray, D., Wade, B., 2018. The Evolution of the Damara Orogenic System: A Record of West Gondwana Assembly and Crustal Response. *Geology of the SW Gondwana*. In: Siegesmund, S., Oyhantçabal, P. and Basei, M.A.S., Oriolo, S. (Eds.), *Regional Geology Review*, Springer, pg. 303-386.
- Harrison, T.M., Blichert-Toft, J., Muller, W., Albarede, F., Holden, P.; Mojzsis, S.J., 2005. Heterogeneous Hadean hafnium: evidence for continental crust at 4.4 to 4.5 Ga. *Science*, v.310, p.1947.1950.
- Hanchar, J.M. & Miller, C.F., 1993. Zircon zonation patterns as revealed by cathodoluminescence and backscattered electron images: Implications for interpretation of complex crustal histories. *Chemical Geology*. 100: 1-13.
- Hawkesworth, C., Cawood, P., Kemp, T., Storey, C., and Dhuime, B., 2009. A Matter of Preservation: *Science*, v. 323, 49–50, doi:10.1126/science.1168549.
- Hawkesworth, C., Dhuime, B., Pietranik, A., Cawood, P., Kemp, T., and Storey, C., 2010. The Generation and Evolution of the Continental Crust: *Journal of the Geological Society*, v. 167, p. 229–248, doi:10.1144/0016-76492009-072.
- Hueck, M., Oyhantçabal, P., Philipp, R.P., Basei, M.A.S., Siegesmund, S., 2018. The Dom Feliciano Belt in Southern Brazil and Uruguay. *Geology of the SW Gondwana*. In: Siegesmund, S., Oyhantçabal, P. and Basei, M.A.S., Oriolo, S. (Eds.), *Regional Geology Review*, Springer, pg. 267-303.

- Hueck, M., Oriolo, S., Basei, M.A.S., Oyhantçabal, P., Heller, B.M., Wemmer K., Siegesmund, S., 2022. Archean to early Neoproterozoic crustal growth of the southern South American Platform and its wide-reaching “African” origins. *Precambrian Res.* 369:106532.
- Iglesias, C. M. Da F.; Camozzato, E.; Klein, C. Carta geológica Curral de Pedras, Folha SI.22-V-A-I, estado do Rio Grande do Sul. Porto Alegre: CPRM, 2018. Escala 1:100.000. Programa Geologia do Brasil – PGB.
- Iglesias, C. M. Da F.; Klein, C.; Camozzato, E. Carta geológica Passo São Diogo, Folha SH.22-Y-C-IV. Porto Alegre: CPRM, 2018. Escala 1:100.000. Programa Geologia do Brasil - PGB.
- Janoušek, V., Farrow, C.M., Erban, V., 2006. Interpretation of whole-rock geochemical data in igneous geochemistry: Introducing Geochemical Data Toolkit (GCDkit). *Journal of Petrology* 47, 1255–1259.
- Klein, F.G., Koester, E., Vieira, D.T., Porcher, C.C., Ramos, R.C., Philipp, R.P., 2018. Geologia do Granito Três Figueiras: magmatismo peraluminoso de 585 Ma no sudeste do Cinturão Dom Feliciano. *Pesqui. em Geociências* 45 (2), e0665.
- Loureiro, P.O., Koester, E., Weinberg, R.F., Lenz, C., Porcher, C.C. Ramos, R.C., Vieira, D.T., Bastos, V.A., Knijnik, D.B., Pimentel, M.M. 2021. Recycling and hybridization of incremental episodes of magma intrusions: Pinheiro Machado Complex, Southeastern Dom Feliciano Belt, Brazil. *J. S. Am. Earth Sci.* 105, 102922.
- Ludwig, K. R., 1998. Using Isoplot/Es (Version 1.00b) A Geochronological Toolkit for Microsoft Excel. Berkley Geochronology Center, Special Publication N°1 44p.
- Konopásek, J., Cavalcante, C., Fossen, H., Janousek, V., 2020. Adamastor – an ocean that never existed? *Earth Sci. Rev.* 205, 103201.
- Meira, V.T., García-Casco A., Juliani C., Almeida R.P., Hans, J, Schorscher D., 2015. The role of intracontinental deformation in supercontinent assembly: insights from the

- Ribeira Belt, Southeastern Brazil (Neoproterozoic West Gondwana). *Terra Nova*, 0(0): 1-12.
- Parrish, R. & Noble, S., 2003. Zircon U-Th-Pb Geochronology by Isotope Dilution-Thermal Ionization Mass Spectrometry (ID-TIMS), in J. Hanchar and P. Hoskin (eds) *Zircon: Reviews of Mineralogy and Geochemistry*, Mineralogical Society of America, 183-213.
- Percival, J.J., Konopásek, J., Eiesland, R., Sláma, J., Campos, R.C., Battisti, M.A., Bitencourt, M.F., 2021. Pre-orogenic connection of the foreland domains of the Kaoko–Dom Feliciano–Gariép orogenic system. *Precambrian Res.* 354, 106060.
- Philipp R.P., Machado R., Nardi L.V.S., Lafon J.M. 2002. O magmatismo granítico Neoproterozóico do Batólito Pelotas no sul do Brasil: novos dados e revisão de geocronologia regional. *Revista Brasileira de Geociências*, 32(2):277-290.
- Philipp, R.P., & Machado, R., 2005. The Late Neoproterozoic granitoid magmatism of the Pelotas Batholith, southern Brazil. *J. S. Am. Earth Sci.* 19, 461-478.
- Philipp, R. P., & Machado, R., 2005. The Late Neoproterozoic granitoid magmatism of the Pelotas Batholith, southern Brazil. *Journal of South American Earth Sciences*.
- Philipp, R.P., Pimentel, M.M., Chemale Jr, F., 2016. Tectonic evolution of the Dom Feliciano Belt in Southern Brazil: Geological relationships and U-Pb geochronology. *Braz. J. Geol.* 46 (1), 83-104.
- Oriolo, S., Oyhantçabal, P., Konopásek, J., Basei, M.A.S., Frei, R., Slama, J., Wemmer, K., Siegesmund, S., 2019. Late paleoproterozoic and mesoproterozoic magmatism of the Nico Pérez terrane (Uruguay): tightening up correlations in southwestern Gondwana. *Precambrian Res.* 327, 296–313.
- Oyhantçabal, P., Siegesmund, S., Wemmer, K., 2011. The Río de la Plata Craton: a review of units, boundaries, ages, and isotopic signature. *Int. J. Earth Sci.* 100, 201–220.

- Porada, H., 1989. Pan-African rifting and orogenesis in southern to equatorial Africa and eastern Brazil. *Precambr Res* 44:103–136.
- Preciozzi, F., Masquelin, H., Basei, M.A.S., 1999. The Namaqua/Grenville Terrane of eastern Uruguay. *South America Symposium on Isotope Geology, Actas 58*. Cordoba, Argentina, pp 338–340.
- Ramos, R. C., & Koester, E. (2014). Geology of the Arroio Grande Metamafic-Ultramafic Association, Southeastern Sul-Rio-Grandense Shield. *Pesquisas Em Geociencias*.
- Rubatto, D., Gebauer, D., 1998. Use of cathodoluminescence for U–PB zircon dating by ion microprobe: some examples from the Western Alps. In: Pagel, M., Barbin, V., Blanc, P., Ohnenstetter, D. (Eds.), *Cathodoluminescence in Geosciences*. Springer-Verlag, Berlin, pp. 373–400.
- Rutherford, E. 1905. Present problem in radio-activity. *Popular Science Monthly*, 67: 5-34.
- Scholl, D.W., von Huene, R., 2009. Implications of estimated magmatic additions and recycling losses at the subduction zones of accretionary (non-collisional) and collisional (suturing) orogens. In: Cawood, P.A., Kröner, A. (Eds.), *Earth Accretionary Systems in Space and Time*. Special Publication 318. Geological Society, London, London, pp. 105–125
- Silva, L.C., McNaughton, N. J., Armstrong, R., Hartmann, L. A., Fletcher, I. R., 2005. The Neoproterozoic Mantiqueira Province and its African connections, a zircon-based U–Pb geochronologic subdivision of the Brasiliano/Pan-African systems of orogens. *Precamb. Res.* 136, 203-240.
- Silva, R. F. O granito Capão do Leão: magmatismo tipo-I altamente fracionado no sudeste do Cinturão Dom Feliciano, RS. 2016. 75 f. Master Dissertation-Universidade Federal do Rio Grande do Sul, Porto Alegre, 2016.
- Silva Lara, H.S., Siegesmund, S., Oriolo, S., Hueck, M., Wemmer, K., Basei, M.A.S., Oyhantçabal, P., 2022. Reassessing the polyphase Neoproterozoic evolution of the

Punta del Este Terrane, Dom Feliciano Belt, Uruguay. *Int. J. Geol.*
doi.org/10.1007/s00531-022-02230-0.

Storey, B.C., 1995, The role of mantle plumes in continental break-up: case histories from Gondwanaland: *Nature*, v. 377, p. 301–308, doi:10.1038/377301a0.

Takehara, L.; Laux, J.H., 2019. Área de Relevante Interesse Mineral Integração geológica-geofísica e recursos minerais Batólito Pelotas e Terreno Tijudas; Escala 1:500.000/Lucy Takehara, Jorge Henrique Laux. – Porto Alegre: CPRM.

Torquato, J.R., Cordani, 1981. Brazil-Africa geological links. *Earth Science Reviews* 17, 155–176.

Vermeesch, P., 2004. How many grains are needed for a provenance study? *Earth and Planetary Science Letters* 224, 441– 451

Vieira, D.T., Koester, E., Bertotti, A.L., 2016. Petrologia do granito Chasqueiro, região de Arroio grande, sudeste do Escudo sul-rio-grandense. *Braz. J. Geol.* 46, 79–108.

Wetherill G.W., 1956. Discordant uranium-lead ages, I. *Transactions of the American Geophysical Union* 37: 320-326.

Wendt, I., 1984. A tree-dimensional U-Pb Discordia plane to evaluate samples with common lead of unknown isotopic composition. *Isotopic Geoscience* 2:1-12.

White, R.S., 1992. Crustal structure and magmatism of North Atlantic continental margins: *Journal of the Geological Society*, v. 149, p. 841–854.