THE GEOLOGY OF THE GOLD MINERALIZATIONS IN TAPAJÓS MINERAL PROVINCE, AMAZONIAN AREA, BRAZIL

Maria Glícia da Nóbrega Coutinho¹, François Robert ² and Reginaldo Alves dos Santos ¹

¹ Geological Survey of Brazil – CPRM, Av. Pasteur, 404, Rio de Janeiro, Brasil, 22292 170 2 Geological Survey of Canada – 601 Booth Street, Ottawa, Ontario, Canada, K1A OE8

Keywords: Amazonian area, Tapajós geology, Gold mineralizations, Lode-gold deposit, Porphyry-gold, Brazil

Abstract: The geology of the gold mineralizations in Tapajós Mineral Province, Amazonian area, Brazil. The Precambrian Tapajós Mineral Province (90,000 km²) located on the Amazonian Craton, more specifically on Brazil Central Shield underwent a tectonic evolution at about 2.0 Ga to that of the Guiana Shield (Rogers 1996), which has been important to the gold mineralization (Robert 1996). Gold was first discovered in the province in the early 1950's and the exploitation by garimpeiros has continued for 50 years. The cumulative production is estimated to be 159 tonnes of Au from 1958 to 1966. However, no research has been yet attempted to understanding the geology of the gold mineralization. There is thus an urgent need to understand the geology of the these gold deposits, so as to enable an efficient exploration philosophy to be developed.

The province consists of metamorphic complex basement (2.016 Ga and 2.44 Ga; Santos et al. 1997), granitic-gneissic migmatitic, cut by NWstrike-slip shear zone associated compressional regime, predominantly ductile (Santos 1996). Supracrustal rocks dated at 2.1 Ga (Santos et al. 1997) are mainly schist, metachert, subordinated amphibolite. Mafic rocks (1.85 Ga; Santos et al. 1997) and granitoids (1.90 Ga) cut the basement. A expressive calc-alkaline magmatic event (Gibbs and Baron 1993) consisting of felsic volcanics (1.82 Ga) and subvolcanic rocks (1.77 Ga) is marked by extensional brittle regime (Santos Sedimentary cover correlated to Roraima Supergroup is intruded by basic dykes and sills (1.69 Ga; Gibbs and Baron 1993). Mafic intrusions dated at 1.2 and 1.0 Ga also occur. Cenozoic sedimentary cover form the alluvial deposits and are gold-rich.

Two type of gold mineralizations occurs: (i) gold-bearing quartz veins (lode-gold deposits) close associated with calc-alkaline magmatism, genetically linked to subduction; and (ii) disseminated/stockwork

mineralizations (porphyry-gold) associated with calcalkaline to alkaline magmatism during an extensional regime at a high crustal level (Robert 1996; Coutinho *et al.* 1997, 1998).

REFERENCES

Gibbs, A.K. & C.N. Barron, 1993. The geology of the Guiana Shield. *Oxford Monog. on Geology & Geophysics*, **22**, Oxford University, New York, 246 p Coutinho, M.G. da N., F. Robert and R.A. dos Santos, 1997. A first approach in understanding the gold mineralization of the Tapajós Mineral Province, Amazonian area, *Brazil. Gold'97*, 10th, Rio de Janeiro, Brazil.

Coutinho, M.G. da N., T. Liverton, E.C. de Souza, 1998. Granitic magmatism and related gold mineralizations in Tapajós Mineral Province, Amazonian area, Brazil. *Journal of South America Earth Sciences* (in press).

Robert, F., 1996. Tapajós Gold Project Pará State, Brazil. *Mission Report, CIDA Project* 204/13886, Canada-Brazil Cooperation Project, 35 p.

Rogers, J.J.W., 1996. History of continents in the past three billion years. *Journal of Geology*, 109, p 91-107.

Santos, J.O.S., L.A. Hartmann and H.E. Gaudette, 1997. Reconnaissance U/Pb in zircon, Pb/Pb in sulfides and review of Rb/Sr geochronology in the Tapajós gold Province, Brazil. *South America Symposium on Isotopo Geology*, Campos de Jordão, SP, p 280-282.

Santos, R.A. dos, 1996. Contribuição à análise estrutural de jazimentos auríferos do rio Tapajós, SW do Pará - Garimpos: Chico Torres, São José, Mamoal, Jutaí, Abacaxis, Espírito Santo, Bom Jesus, Goiano, Fazenda Pizon, Ouro Mil, Santa Isabel, Majestade, Carneirinho e Creporizão. Relotório Internos, vols. I, II & III, Companhia de Pesquisa de Recursos Minerais, Salvador, Brasil, 81 p.